

The bicanonical map of surfaces with $p_g = 0$ and $K^2 \geq 7$ *

Margarida Mendes Lopes

Rita Pardini

Abstract

A minimal surface of general type with $p_g(S) = 0$ satisfies $1 \leq K^2 \leq 9$ and it is known that the image of the bicanonical map φ is a surface for $K_S^2 \geq 2$, whilst for $K_S^2 \geq 5$, the bicanonical map is always a morphism. In this paper it is shown that φ is birational if $K_S^2 = 9$ and that the degree of φ is at most 2 if $K_S^2 = 7$ or $K_S^2 = 8$.

By presenting two examples of surfaces S with $K_S^2 = 7$ and 8 and bicanonical map of degree 2, it is also shown that this result is sharp. The example with $K_S^2 = 8$ is, to our knowledge, a new example of a surface of general type with $p_g = 0$.

The degree of φ is also calculated for two other known surfaces of general type with $p_g = 0$, $K_S^2 = 8$. In both cases the bicanonical map turns out to be birational.

1 Introduction

Many examples of complex surfaces of general type with $p_g = q = 0$ are known, but a detailed classification is still lacking, despite much progress in the theory of algebraic surfaces. Surfaces of general type are often studied using properties of their canonical curves. If a surface has $p_g = 0$, then there are of course no such curves, and it seems natural to look instead at the bicanonical system, which is not empty.

Minimal surfaces S of general type with $p_g(S) = 0$ satisfy $1 \leq K_S^2 \leq 9$. By a result of Xiao Gang [15], the image of the bicanonical map is a surface

*2000 Mathematics Subject Classification: 14J29, 14E05

for $K_S^2 \geq 2$, while for $K_S^2 \geq 5$, by Reider's theorem [14], the bicanonical map is always a morphism. Xiao Gang [16] showed that the degree of the bicanonical map is ≤ 2 for surfaces of general type, with a limited number of possible exceptions. Surfaces with $p_g = 0$ are among the exceptional cases, and [16] gives practically no information on the possible degrees of their bicanonical maps.

The first author [9] showed that if $K_S^2 \geq 3$ and the bicanonical map is a morphism, its degree is ≤ 4 . There are examples due to Burniat [3] (see also [13]) with $3 \leq K^2 \leq 6$ having bicanonical map of degree 4; indeed, [10] gives a precise description of the surfaces with $K_S^2 = 6$, $p_g(S) = 0$ and bicanonical map of degree 4. Here we refine the result of [9] by proving the following results:

Theorem 1.1 *Let S be a minimal surface of general type defined over \mathbb{C} with $p_g(S) = 0$, and $\varphi: S \rightarrow \Sigma \subset \mathbb{P}^{K_S^2}$ its bicanonical map, with image Σ .*

- (i) *If $K_S^2 = 9$ then φ is birational;*
- (ii) *if $K_S^2 = 7, 8$ then φ has degree ≤ 2 .*

Proposition 1.2 *There exist minimal surfaces of general type with $p_g(S) = 0$, $K_S^2 = 7, 8$ and bicanonical map of degree 2.*

We prove this by giving two examples of surfaces S with $K_S^2 = 7$ and 8 and bicanonical map of degree 2. The example with $K_S^2 = 7$ is due to Inoue [8, Remark 6], who constructed it as a quotient of a complete intersection in the product of four elliptic curves by a free action of \mathbb{Z}_2^5 . Here we give an alternative description as a \mathbb{Z}_2^2 -cover of a singular rational surface that allows us to describe the bicanonical map and compute its degree. The example with $K_S^2 = 8$ is obtained by applying a construction of Beauville ([1, p. 123, Exercise 4], and cf. [7]). To the best of our knowledge, it is a new example of a surface of general type with $p_g = 0$.

Only a few examples of surfaces with $p_g = 0$, $K_S^2 = 8$ are known. It is a nontrivial exercise to compute the degree of the bicanonical map for an explicit surface, and for interest, we include the computation for two other known surfaces of general type with $p_g = 0$, $K_S^2 = 8$, for both of which the bicanonical map turns out to be birational.

The paper is organized as follows: Section 2 recalls some facts on irregular double covers, one of the main ingredients of the proof of the theorem; in

Section 3 we prove the main result. For $K_S^2 = 7, 8$ the proof consists of using the methods of Section 2 to exclude the possibility that the bicanonical map has degree 4; for $K_S^2 = 9$ the result is obtained by combining Reider's theorem with an analysis of the Picard group of S . In the final Section 4 we present the two examples to prove Proposition 1.2, and we also compute the degree of the bicanonical map of two other surfaces with $p_g = 0$ and $K_S^2 = 8$.

Notations and conventions We work over \mathbb{C} ; all varieties are assumed to be compact and algebraic. We do not distinguish between line bundles and divisors on a smooth variety, and use additive and multiplicative notation interchangeably. Linear equivalence is denoted by \equiv and numerical equivalence by \sim . The remaining notation is standard in algebraic geometry.

2 Irregular double covers and fibrations

We describe here the key facts used in some proofs in this paper.

Let S be a smooth complex surface, $D \subset S$ a curve (possibly empty) with at worst ordinary double points, and M a line bundle on S with $2M \equiv D$. It is well known that there exists a normal surface Y and a finite degree 2 map $\pi: Y \rightarrow S$ branched over D such that $\pi_*\mathcal{O}_Y = \mathcal{O}_S \oplus M^{-1}$. The singularities of Y are A_1 points and occur precisely above the singular points of D ; thus it makes sense to speak of the canonical divisor, the geometric genus, the irregularity and the Albanese map of Y . We refer to Y as the *double cover defined by the relation $2M \equiv D$* . The invariants of Y are:

$$\begin{aligned} K_Y^2 &= 2(K_S + M)^2, \\ \chi(\mathcal{O}_Y) &= 2\chi(\mathcal{O}_S) + \frac{1}{2}M(K_S + M), \\ p_g(Y) &= p_g(S) + h^0(S, K_S + M). \end{aligned} \tag{2.1}$$

If $p_g(S) = q(S) = 0$, the existence of a double cover $\pi: Y \rightarrow S$ with $q(Y) > 0$ forces the existence of a fibration $f: S \rightarrow \mathbb{P}^1$ such that π^{-1} of the general fibre of f is disconnected. More precisely we have:

Proposition 2.1 (De Franchis) *Let S be a smooth surface with $p_g(S) = q(S) = 0$ and $\pi: Y \rightarrow S$ a double cover with at most A_1 points; if $q(Y) > 0$, then*

- (i) the Albanese image of Y is a curve B ;
- (ii) let $\alpha: Y \rightarrow B$ be the Albanese fibration. Then there exists a fibration $g: S \rightarrow \mathbb{P}^1$ and a degree 2 map $p: B \rightarrow \mathbb{P}^1$ such that $p \circ \alpha = g \circ \pi$.

The possibility of existence of such a double cover often leads to a contradiction, using the following result:

Corollary 2.2 *Let S be a minimal surface of general type with $p_g(S) = q(S) = 0$ and $K_S^2 \geq 3$, and $\pi: Y \rightarrow S$ a double cover with at most A_1 points. Then $K_Y^2 \geq 16(q(Y) - 1)$.*

Proposition 2.1 is an old result of De Franchis [6], explained and generalized in several ways by Catanese and Ciliberto [5]. Proposition 2.1 and Corollary 2.2 are both stated and proved in [10] for smooth Y , but the proof extends verbatim to the case of A_1 points.

3 Proof of Theorem 1.1

Under the assumptions of Theorem 1.1, the image of the bicanonical map is a surface by [15], and the bicanonical map is a morphism by Reider's theorem [14]. Moreover, since $4K_S^2 = \deg \varphi \deg \Sigma$ and Σ is a nondegenerate surface in $\mathbb{P}^{K_S^2}$, the possible values of $\deg \varphi$ are 1, 2, 4 for $K_S^2 = 7, 8$ and 1, 2, 3, 4 for $K_S^2 = 9$.

We prove the theorem by analysing separately the cases $K_S^2 = 7, 8, 9$. In each case we argue by contradiction.

3.1 The case $K_S^2 = 7$

By the above remark, it is enough to show that $\deg \varphi = 4$ does not occur. Assume that φ has degree 4. The bicanonical image Σ is a linearly normal surface of degree 7 in \mathbb{P}^7 and its nonsingular model has $p_g = q = 0$. By [11, Theorem 8], Σ is the image of the blowup $\widehat{\mathbb{P}}$ of \mathbb{P}^2 at two points P_1, P_2 under its anticanonical map $f: \widehat{\mathbb{P}} \hookrightarrow \mathbb{P}^7$. If $P_1 \neq P_2$, then f is an embedding, while if P_2 is infinitely near to P_1 (say) then Σ has an A_1 singularity. In either case, the hyperplane section of Σ can be written as $H \equiv 2l + l_0$, where l is the image on Σ of a general line of \mathbb{P}^2 and l_0 is the image on Σ of the strict transform of the line through P_1 and P_2 . Notice that l_0 is contained

in the smooth part of Σ . Thus we have $2K_S \equiv 2L + L_0$, where $L = \varphi^*l$ and $L_0 = \varphi^*l_0$.

Lemma 3.1 *L_0 satisfies one of the following possibilities:*

- (i) *there exists an effective divisor D on S such that $L_0 = 2D$; or*
- (ii) *L_0 is a smooth rational curve with $L_0^2 = -4$; or*
- (iii) *there exist smooth rational curves A and B with $A^2 = B^2 = -3$, $AB = 1$, and $L_0 = A + B$.*

Proof Remark first that $K_S L_0 = 2$, $L_0^2 = -4$, and $L_0 = 2(K_S - L)$ is divisible by 2 in $\text{Pic } S$. Let θ be a -2 -curve of S ; then θ is contracted by φ and thus $L\theta = L_0\theta = 0$. Since L and L_0 are independent elements of the 3-dimensional space $H^{1,1}(S)$, S contains at most one -2 -curve. We write $L_0 = C + a\theta$, where C is the strict transform of L_0 , θ is a -2 -curve and $a \geq 0$ (we set $a = 0$ if S has no -2 -curve). The equalities $\theta L_0 = 0$ and $L_0^2 = -4$ imply

$$\theta C = 2a, \quad \text{and} \quad C^2 = -4 - 2a^2. \quad (3.1)$$

If C is irreducible, then $K_S C = 2$ implies $C^2 \geq -4$ and thus $a = 0$ and case (ii) holds. If C is reducible, then $C = A + B$, with A and B irreducible curves such that $K_S A = K_S B = 1$. If $A = B$, then $AL_0 = 2A^2 + a\theta A = 2A^2 + a^2$ is even, because L_0 is divisible by 2, and thus a is even and we are in case (i). If $A \neq B$, then $AB \geq 0$ and $A^2, B^2 \geq -3$; by parity considerations and (3.1) we get $A^2 = B^2 = -3$ and either $AB = 1$, $a = 0$ or $AB = 0$, $a = 1$. The first case corresponds to (iii), while the second does not occur. In fact the intersection matrix of A, B, θ would be negative definite, contradicting the index theorem, since $h^{1,1}(S) = 3$. \diamond

In cases (ii) or (iii) of Lemma 3.1, let $\pi: Y \rightarrow S$ be the double cover given by $2(K_S - L) \equiv L_0$; then the formulas (2.1) give $\chi(Y) = 2$ and $K_Y^2 = 16$. Since the bicanonical map φ maps L onto a twisted cubic, $h^0(S, \mathcal{O}_S(2K_S - L)) = 4$ and thus $p_g(Y) = p_g(S) + h^0(S, \mathcal{O}_S(2K_S - L)) = 4$; we thus obtain $q(Y) = 3$, contradicting Corollary 2.2.

In case (i) of Lemma 3.1, consider the étale double cover $\pi: Y \rightarrow S$ given by $2(K_S - L - D) \equiv 0$; arguing as above, we get that the invariants of Y are

$$K_Y^2 = 14, \quad \chi(\mathcal{O}_Y) = 2, \quad p_g(Y) = p_g(S) + h^0(S, \mathcal{O}_S(2K_S - L - D)) = 3,$$

so that $q(Y) = 2$ and we again obtain a contradiction to Corollary 2.2.

Hence $\deg \varphi \neq 4$ and we have proved Theorem 1.1 in case $K_S^2 = 7$.

3.2 The case $K_S^2 = 8$

As in case $K_S^2 = 7$, it is enough to show that $\deg \varphi = 4$ does not occur. If φ has degree 4, then the bicanonical image Σ is a linearly normal surface of degree 8 in \mathbb{P}^8 whose nonsingular model has $p_g = q = 0$. By [11, Theorem 8], Σ is either the Veronese embedding in \mathbb{P}^8 of a quadric $Q \subset \mathbb{P}^3$ or the image of the blowup $\widehat{\mathbb{P}}$ of \mathbb{P}^2 at a point P under its anticanonical map $f: \widehat{\mathbb{P}} \hookrightarrow \mathbb{P}^8$.

In the first case $2K_S \equiv 2A$, where A is the hyperplane section of Q . Then $\eta = K_S - A$ is a nontrivial 2-torsion element in $\text{Pic } S$, since $p_g(S) = 0$. The étale double cover $\pi: Y \rightarrow S$ given by $2\eta \equiv 0$ has invariants $\chi(Y) = 2$, $K_Y^2 = 16$. Moreover, $p_g(Y) = p_g(S) + h^0(S, \mathcal{O}_S(A)) = 4$, so that $q(Y) = 3$. Since $K_Y^2 = 16$, this contradicts Corollary 2.2, and therefore Σ is not the Veronese embedding of a quadric.

If the bicanonical image Σ is the image of $\widehat{\mathbb{P}}$ via the map induced by $|-K_{\widehat{\mathbb{P}}}|$, then the hyperplane section of Σ can be written as $H \equiv 2l + l_0$, where l is the image on Σ of a general line of \mathbb{P}^2 and l_0 is the image on Σ of the strict transform of a general line through P . Thus $2K_S \equiv 2L + L_0$, where $L = \varphi^*l$ and $L_0 = \varphi^*l_0$, and $L_0 = \varphi^*l_0$ is smooth by Bertini's theorem. Consider now the double cover $\pi: Y \rightarrow S$ given by $2(K_S - L) \equiv L_0$; the formulas (2.1) give $\chi(Y) = 3$ and $K_Y^2 = 24$. Since $p_g(Y) = p_g(S) + h^0(S, \mathcal{O}_S(2K_S - L)) = 0 + h^0(S, \mathcal{O}_S(L + L_0)) = 5$, we get $q(Y) = 3$, contradicting Corollary 2.2. Thus Σ is also not the image of $\widehat{\mathbb{P}}$.

Hence $\deg \varphi \neq 4$ and the proof of Theorem 1.1, (ii) is complete.

3.3 The case $K_S^2 = 9$

If $K_S^2 = 9$, then by Poincaré duality, $H^2(S, \mathbb{Z})$ is generated up to torsion by the class of a line bundle L with $L^2 = 1$; thus every divisor on S is numerically a multiple of L , and in particular $K_S \sim 3L$.

Assume by contradiction that φ is not birational; then by Reider's theorem (cf. [2, Theorem 2.1]), for every pair of points $x_1, x_2 \in S$ with $\varphi(x_1) = \varphi(x_2)$ there exists an effective divisor C containing x_1, x_2 such that $K_S C - 2 \leq C^2 < \frac{1}{2} K_S C < 2$. Since $K_S \sim 3L$, the only possibility is that $C \sim L$. We can assume that, as x_1 and x_2 vary, the divisor C varies in an irreducible system of curves, which is linear by the regularity of S . Every curve of $|C|$

is irreducible, since the class of C generates $H^2(S, \mathbb{Z})$ up to torsion, and the general curve of $|C|$ is smooth by Bertini's theorem, since $C^2 = 1$. Therefore $|C|$ is a linear pencil of curves of genus 3 with one base point. For a general $C \in |C|$ we consider the exact sequence:

$$0 \rightarrow \mathcal{O}_S(2K_S - C) \rightarrow \mathcal{O}_S(2K_S) \rightarrow \mathcal{O}_C(2K_S) \rightarrow 0. \quad (3.2)$$

Since $2K_S - C \sim K_S + 2L$, Kodaira vanishing gives $H^1(S, \mathcal{O}_S(2K_S - C)) = 0$, and the map $H^0(S, \mathcal{O}_S(2K_S)) \rightarrow H^0(C, \mathcal{O}_C(2K_S))$ induced by the sequence (3.2) is surjective. So the map $f: C \rightarrow \mathbb{P}^3$ given by $|\mathcal{O}_C(2K_S)|$ is not birational; it follows that f maps C two-to-one onto a twisted cubic, and thus C is hyperelliptic. If we denote by Δ the g_2^1 of C , then $2K_S|_C \equiv 3\Delta$ and also, by the adjunction formula, $K_S + C|_C \equiv 2\Delta$. So $\eta \equiv 4K_S - (3K_S + 3C) \equiv K_S - 3C$ is trivial when restricted to C . Moreover $\eta \sim 0$ and so η is a torsion element of $\text{Pic } S$. Since $p_g(S) = 0$, η is nonzero. Consider the connected étale cover $\pi: Y \rightarrow S$ associated to η . Because $\eta|_C = 0$, the cover $\pi|_{\pi^{-1}(C)}: \pi^{-1}(C) \rightarrow C$ is trivial and thus $\pi^{-1}(C)$ is a smooth disconnected curve with each component of self-intersection 1. This contradicts the Index theorem and we have thus proved Theorem 1.1, (i). \diamond

4 Examples


This section calculates the degree of the bicanonical map in 4 interesting examples, as discussed in the introduction.

Example 4.1 Starting from the quadrilateral $P_1P_2P_3P_4$ in \mathbb{P}^2 of Figure 1, let P_5 be the intersection point of the lines P_1P_2 and P_3P_4 and P_6 the intersection point of P_1P_4 and P_2P_3 . Write $\Sigma \rightarrow \mathbb{P}^2$ for the blowup of P_1, \dots, P_6 , and e_i for the exceptional curves of Σ over P_i . Denote by l the pullback of a line.

Write S_1, \dots, S_4 for the strict transforms on Σ of the sides P_iP_{i+1} of the quadrilateral $P_1P_2P_3P_4$ (we take subscripts modulo 4); these are the only -2 -curves of Σ . The morphism $f: \Sigma \rightarrow \mathbb{P}^3$ given by $|-K_\Sigma|$ has image a cubic surface $V \subset \mathbb{P}^3$, and f is an isomorphism on $\Sigma \setminus \bigcup S_i$, and contracts each S_i to an A_1 point.

If $A \subset \{P_1, \dots, P_6\}$ consists of 4 points no three of which are collinear, then the linear system of conics through the points of A gives rise to a free pencil on Σ ; we denote by f_1 the strict transform of a general conic through

Finally, we introduce the “diagonals” of the quadrilateral $P_1P_2P_3P_4$, writing $\Delta_1, \Delta_2, \Delta_3$ for the strict transform of P_1P_3 , P_2P_4 and P_5P_6 . The divisors



we have introduced satisfy the following relations:

- Denote by $\gamma_1, \gamma_2, \gamma_3$ the nonzero elements of $\Gamma = \mathbb{Z}_2 \times \mathbb{Z}_2$ and by $\chi_i \in \Gamma^*$ the nontrivial character orthogonal to γ_i ; by [12, Propositions 2.1 and 3.1], to define a smooth Γ -cover $\pi: X \rightarrow \Sigma$, we specify:

- 8

The branch locus of π is D . More precisely, D_i is the image of the divisorial part of the fixed locus of γ_i on S . We have

$$\pi_* \mathcal{O}_S = \mathcal{O}_\Sigma \oplus L_1^{-1} \oplus L_2^{-1} \oplus L_3^{-1},$$

where $L_3 = L_1 + L_2 - D_3$, and Γ acts on L_i^{-1} via the character χ_i .

Here we set:

(I) $D_1 = \Delta_1 + f_2 + S_1 + S_2$, $D_2 = \Delta_2 + f_3$, $D_3 = \Delta_3 + f_1 + f'_1 + S_3 + S_4$;
where $f_1, f'_1 \in |f_1|$, $f_2 \in |f_2|$, $f_3 \in |f_3|$ are general curves;

(II) $L_1 = 5l - e_1 - 2e_2 - e_3 - 3e_4 - 2e_5 - 2e_6$, and

$$L_2 = 6l - 2e_1 - 2e_2 - 2e_3 - 2e_4 - 3e_5 - 3e_6$$

and we obtain $L_3 = 4l - 2e_1 - 2e_2 - 2e_3 - e_4 - e_5 - e_6$. For $i = 1, \dots, 4$, the (set theoretic) inverse image of S_i in X is the disjoint union of two -1 -curves E_{i1}, E_{i2} ; contracting these 8 exceptional curves on X and contracting the S_i on Σ , we obtain a smooth \mathbb{Z}_2^2 -cover $p: S \rightarrow V$. The map p is branched on the four singular points of V and on the image \overline{D} of D , which is contained in the smooth locus of V . The bicanonical divisor $2K_X$ is equal to $\pi^*(2K_\Sigma + D) = \pi^*(-K_\Sigma + f_1 + S_1 + S_2 + S_3 + S_4) = \pi^*(-K_\Sigma + f_1) + 2\sum E_{ij}$, and thus the bicanonical divisor $2K_S$ is equal to $\pi^*(-K_V + \overline{f}_1)$, where \overline{f}_1 is the image of f_1 in V . So $2K_S$ is ample, since it is the pullback of an ample line bundle by a finite map, S is minimal and of general type, and $K_S^2 = \frac{1}{4}4(K_V + \overline{f}_1)^2 = 7$.

To compute the geometric genus of S , recall that $p_g(X) = p_g(\Sigma) + \sum h^0(\Sigma, K_\Sigma + L_i)$ (cf. [4] or [12, Lemma 4.2]). We have

$$\begin{aligned} K_\Sigma + L_1 &= 2l - e_2 - 2e_4 - e_5 - e_6, \\ K_\Sigma + L_2 &= 3l - e_1 - e_2 - e_3 - e_4 - 2e_5 - 2e_6, \\ K_\Sigma + L_3 &= l - e_1 - e_2 - e_3. \end{aligned}$$

We show that $h^0(\Sigma, K_\Sigma + L_2) = 0$. Assume by contradiction that there exists $D \in |K_\Sigma + L_2|$ and consider the image C of D in \mathbb{P}^2 ; C is a cubic containing P_1, \dots, P_6 which has a double point at P_5 and P_6 . By Bezout's theorem, Δ_3 is contained in C and thus $C = \Delta_3 + Q$, where Q is a conic containing P_1, \dots, P_6 , which is impossible. By similar (easier) arguments, one shows that $h^0(\Sigma, K_\Sigma + L_1) = h^0(\Sigma, K_\Sigma + L_3) = 0$, and thus $p_g(S) = p_g(X) = 0$. By the projection formula for a finite flat morphism,

$$H^0(X, 2K_X) =$$

$$H^0(\Sigma, -K_\Sigma + f_1 + \sum S_j) \oplus \left(\bigoplus_i H^0(\Sigma, -K_\Sigma + f_1 + \sum S_j - L_i) \right),$$

and Γ acts on $H^0(\Sigma, -K_\Sigma + f_1 + \sum S_j - L_i)$ via the character χ_i . We have $h^0(\Sigma, -K_\Sigma + f_1 + \sum S_j) = h^0(\Sigma, -K_\Sigma + f_1)$, since

$$S_j(-K_\Sigma + f_1 + \sum S_i) = -2 \quad \text{for } i = 1, \dots, 4;$$

in addition, $h^0(\Sigma, -K_\Sigma + f_1) = 7$, since Σ is rational, $2f_1 + f_2 + f_3$ has arithmetic genus 7, and $-K_\Sigma + f_1 = K_\Sigma + 2f_1 + f_2 + f_3$. Since $p_2(S) = 8$, there is a value $i \in \{1, 2, 3\}$ such that $h^0(\Sigma, -K_\Sigma + \sum S_j + f_1 - L_i) = 1$ and $h^0(\Sigma, -K_\Sigma + \sum S_j + f_1 - L_k) = 0$ for $k \neq i$. Actually, an argument similar to that used for computing $p_g(S)$ shows that

$$h^0(-K_\Sigma + \sum S_j + f_1 - L_1) = h^0(\sum S_j + e_4) = 1,$$

$$h^0(-K_\Sigma + \sum S_j + f_1 - L_2) = h^0(3l - e_1 - 2e_2 - e_3 - 2e_4 - e_5 - e_6) = 0,$$

$$h^0(-K_\Sigma + \sum S_j + f_1 - L_3) = h^0(5l - e_1 - 2e_2 - e_3 - 3e_4 - 3e_5 - 3e_6) = 0.$$

It follows that the bicanonical map $\varphi: S \rightarrow \mathbb{P}^7$ is composed with the involution γ_1 but not with γ_2 and γ_3 . Since $|2K_S| \supset \pi^*|-K_\Sigma|$ and the map $\Sigma \rightarrow \mathbb{P}^3$ induced by $|-K_\Sigma|$ is birational, it follows that φ has degree 2.

The remaining examples are obtained using the following construction due to Beauville (see [1, p. 123, Ex. 4] and cf. [7]). Let C_1, C_2 be curves of genus g_1, g_2 , and assume that a group G of order $(g_1 - 1)(g_2 - 1)$ acts on C_1, C_2 so that C_i/G is isomorphic to \mathbb{P}^1 for $i = 1, 2$; write $p_i: C_i \rightarrow \mathbb{P}^1$ for the projections onto the quotients and $p: C_1 \times C_2 \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$ for the product of p_1 and p_2 . Thus p is a Galois cover with group $G \times G$. Assume in addition that there exists an automorphism $\psi \in \text{Aut } G$ whose graph $\Gamma = \Gamma_\psi \subset G \times G$ acts freely on $C_1 \times C_2$. Then set $S = (C_1 \times C_2)/\Gamma$ and denote by $q: C_1 \times C_2 \rightarrow S$ the quotient map and by $\pi: S \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$ the map induced by p . If G is Abelian, then π is a G -cover. The surface S is minimal and of general type since $C_1 \times C_2$ is minimal of general type and q is étale. Since Γ acts freely, $\chi(\mathcal{O}_{C_1 \times C_2}) = |G|\chi(\mathcal{O}_S)$ and $K_{C_1 \times C_2}^2 = |G|K_S^2$, namely $\chi(\mathcal{O}_S) = 1$, $K_S^2 = 8$. The irregularity $q(S)$ equals the dimension of the Γ -invariant subspace of

$H^0(C_1 \times C_2, \Omega_{C_1 \times C_2}^1) \cong H^0(C_1, \omega_{C_1}) \oplus H^0(C_2, \omega_{C_2})$. Since C_1/G and C_2/G are both rational and ψ is an automorphism, it follows that $q(S) = 0$, and thus $p_g(S) = 0$.

Example 4.2 As far as we know, this is a new example. In this case $G = \mathbb{Z}_2^3$, $g_1 = 5$, $g_2 = 3$. We denote by $\gamma_1, \gamma_2, \gamma_3$ the standard generators of G and by χ_1, χ_2, χ_3 the dual basis of the group of characters G^* . To construct the G -cover $p_i: C_i \rightarrow \mathbb{P}^1$ we have to specify (cf. [12, Propositions 2.1 and 3.1]):

- (i) a divisor D_γ on \mathbb{P}^1 for each nonzero $\gamma \in G$;
- (ii) line bundles L_1, L_2, L_3 on \mathbb{P}^1 satisfying

$$2L_i \equiv \sum_{\gamma} \varepsilon_i(\gamma) D_\gamma, \quad \text{where} \quad \begin{cases} \varepsilon_i(\gamma) = 0 & \text{if } \gamma \in \ker \chi_i, \\ \varepsilon_i(\gamma) = 1 & \text{otherwise.} \end{cases}$$

To construct $p_1: C_1 \rightarrow \mathbb{P}^1$, we choose distinct points $P_1, \dots, P_6 \in \mathbb{P}^1$ and set $D_{\gamma_1} = P_1 + P_2$, $D_{\gamma_2} = P_3 + P_4$, $D_{\gamma_3} = P_5 + P_6$, $D_\gamma = 0$ for $\gamma \neq \gamma_i$, and $L_1 = L_2 = L_3 = \mathcal{O}_{\mathbb{P}^1}(1)$. The curve C_1 is smooth connected of genus 5. To construct $p_2: C_2 \rightarrow \mathbb{P}^1$, we choose distinct points $Q_1, \dots, Q_5 \in \mathbb{P}^1$ and set $D_{\gamma_1} = Q_1$, $D_{\gamma_2} = Q_2$, $D_{\gamma_1 + \gamma_2} = Q_3$, $D_{\gamma_3} = Q_4 + Q_5$, $D_\gamma = 0$ for the remaining nonzero elements of G , and $L_1 = L_2 = L_3 = \mathcal{O}_{\mathbb{P}^1}(1)$. The curve C_2 is smooth connected of genus 3. Define $\psi \in \text{Aut } G$ by

$$\gamma_1 \mapsto \gamma_1 + \gamma_3, \quad \gamma_2 \mapsto \gamma_2 + \gamma_3, \quad \gamma_3 \mapsto \gamma_1 + \gamma_2 + \gamma_3.$$

In the above notation, $\pi: S \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$ is a G -cover and $2K_S = \pi^* \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(2, 1)$. By the projection formula, we have

$$H^0(S, 2K_S) = \bigoplus_{\chi \in \Gamma^\perp} H^0(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(2, 1) \otimes M_\chi^{-1}),$$

where M_χ^{-1} is the eigensheaf of $\pi_* \mathcal{O}_S$ corresponding to $\chi \in \Gamma^\perp \cong G^*$, and $(G \times G)/\Gamma \cong G$ acts on $H^0(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(2, 1) \otimes M_\chi^{-1})$ via χ . The M_χ are line bundles on $\mathbb{P}^1 \times \mathbb{P}^1$ that can be determined using [12, (2.15)]. One checks that $H^0(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(2, 1) \otimes M_\chi^{-1})$ is nonzero only for the elements of Γ^\perp that are orthogonal to $(0, \gamma_3) \in G \times G$. It follows that the bicanonical map of S is composed with the involution induced on S by $(0, \gamma_3)$, and thus it has degree 2 by Theorem 1.1.

Example 4.3 As for Example 4.1, this is due to Inoue [8, p. 317], and arises as the quotient of a complete intersection in the product of 4 elliptic curves by a free group action. Here we give a construction in the style of Beauville as explained above which is more suitable for our purpose. Let $\gamma_1, \dots, \gamma_4$ be a basis of $G = \mathbb{Z}_2^4$ and χ_1, \dots, χ_4 the dual basis of G^* ; set $\gamma_0 = \gamma_1 + \gamma_2 + \gamma_3 + \gamma_4$. We construct C_i as G -covers of \mathbb{P}^1 for $i = 1, 2$. As in Example 4.2, for this, we specify (cf. [12, Propositions 2.1 and 3.1]):

- (i) a divisor D_γ of \mathbb{P}^1 for every nonzero $\gamma \in G$;
- (ii) line bundles L_1, \dots, L_4 of \mathbb{P}^1 satisfying

$$2L_i \equiv \sum_{\gamma} \varepsilon_i(\gamma) D_\gamma, \quad \text{where} \quad \begin{cases} \varepsilon_i(\gamma) = 0 & \text{if } \gamma \in \ker \chi_i \\ \varepsilon_i(\gamma) = 1 & \text{otherwise.} \end{cases}$$

Choose distinct points $P_0, \dots, P_4 \in \mathbb{P}^1$ and set $D_{\gamma_i} = P_i$ for $i = 0, \dots, 4$, $D_\gamma = 0$ if $\gamma \neq \gamma_i$, and $L_i = \mathcal{O}_{\mathbb{P}^1}(1)$ for $i = 1, \dots, 4$. We write $p_1: C_1 \rightarrow \mathbb{P}^1$ for the corresponding G -cover. Then C_1 is a smooth connected curve of genus 5. We construct the curve C_2 in the same way, starting from points $Q_0, \dots, Q_4 \in \mathbb{P}^1$.

Let $\psi \in \text{Aut } G$ be the automorphism:

$$\gamma_1 \mapsto \gamma_1 + \gamma_3, \quad \gamma_2 \mapsto \gamma_2 + \gamma_4, \quad \gamma_3 \mapsto \gamma_1 + \gamma_4, \quad \gamma_4 \mapsto \gamma_1 + \gamma_3 + \gamma_4.$$

In the above notation, $\pi: S \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$ is a G -cover and $2K_S = \pi^* \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(1, 1)$. Since $\mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(1, 1)$ is very ample, it follows that the bicanonical map φ of S is birational if and only if it is not composed with an involution γ of the Galois group G of π . To check that this is indeed the case, we use the projection formula

$$H^0(S, 2K_S) = \bigoplus_{\chi \in \Gamma^\perp} H^0(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(1, 1) \otimes M_\chi^{-1}),$$

where M_χ^{-1} is the eigensheaf of $\pi_* \mathcal{O}_S$ corresponding to $\chi \in \Gamma^\perp \cong G^*$, and $(G \times G)/\Gamma \cong G$ acts on $H^0(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(1, 1) \otimes M_\chi^{-1})$ via χ . The M_χ are line bundles on $\mathbb{P}^1 \times \mathbb{P}^1$ that can be determined using [12, (2.15)]. It turns out that no $\gamma \in G \setminus \{0\}$ acts trivially on $H^0(S, 2K_S)$ and thus φ is birational.

Example 4.4 This example is due to Beauville and appears in [1, p. 123, Ex. 4], where the group action is not described explicitly, and in [7]. The assertion concerning the group action in [7] is not correct, since the group action described is not free.

In this case $g = 6$, $G = \mathbb{Z}_5^2$, and $C_1 = C_2 = \{x^5 + y^5 + z^5 = 0\} \subset \mathbb{P}^2$ is the Fermat quintic. If ε is a primitive 5th root of 1, then $(1, 0) \in G$ acts on C by $(x : y : z) \mapsto (\varepsilon x : y : z)$ and $(0, 1)$ acts by $(x : y : z) \mapsto (x : \varepsilon y : z)$. Let ψ be the automorphism of G taking $(1, 0) \mapsto (1, -1)$ and $(0, 1) \mapsto (1, 2)$.

We compute the degree of the bicanonical map of S by writing down an explicit basis of the Γ -invariant subspace of $H^0(C \times C, 2K_{C \times C})$. Take homogeneous coordinates $(x : y : z; x_1 : y_1 : z_1)$ on $\mathbb{P}^2 \times \mathbb{P}^2 \supset C \times C$; using the fact that a regular 1-form on C is the residue of a rational form $\frac{g(x, y, z)}{x^5 + y^5 + z^5} dx \wedge dy \wedge dz$ for g homogeneous of degree 2, we see that $(a, b) \in G$ acts on bicanonical forms on $C \times C$ by:

$$x^i y^j z^{4-i-j} x_1^\alpha y_1^\beta z_1^{4-\alpha-\beta} \mapsto \varepsilon^l x^i y^j z^{4-i-j} x_1^\alpha y_1^\beta z_1^{4-\alpha-\beta},$$

where $l = a(2 + i + \alpha - \beta) + b(3 + j + \alpha + 2\beta)$

Thus the following is a basis of $H^0(Y, 2K_Y)^{\text{inv}}$:

$$\begin{aligned} & x^4 y_1 z_1^3, \quad y^3 z y_1^2 z_1^2, \quad x y z^2 y_1^3 z_1, \quad x^2 y z x_1 z_1^3, \quad z^4 x_1 y_1^3, \\ & x z^3 x_1^2 z_1^2, \quad x^3 y x_1^2 y_1^2, \quad y^4 x_1^3 z_1, \quad x y^2 z x_1^3 y_1. \end{aligned}$$

The subfield of $\mathbb{C}(S)$ generated by ratios of these monomials is the function field $\mathbb{C}(\Sigma)$ of the bicanonical image Σ of S . The map $\pi : S \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$ identifies $\mathbb{C}(\mathbb{P}^1 \times \mathbb{P}^1)$ with the subfield of $\mathbb{C}(S)$ generated by $x^5 z^{-5}$ and $x_1^5 z_1^{-5}$. The extension $\mathbb{C}(S) \supset \mathbb{C}(\mathbb{P}^1 \times \mathbb{P}^1)$ is Galois with Galois group $G = \mathbb{Z}_5^2$. We observe:

$$x^5 z^{-5} = (x^3 y x_1^2 y_1^2)(x^4 y_1 z_1^3)(x^2 y z x_1 z_1^3)^{-1}(z^4 x_1 y_1^3)^{-1}$$

and

$$x_1^5 z_1^{-5} = (z^4 x_1 y_1^3)^2 (y^3 z y_1^2 z_1^2)(x^3 y x_1^2 y_1^2)^2 (x^2 y z x_1 z_1^3)^{-1} (x y z^2 y_1^3 z_1)^{-4}.$$

It follows that $\mathbb{C}(\Sigma) \supset \mathbb{C}(\mathbb{P}^1 \times \mathbb{P}^1)$. Now one checks that no element of the Galois group $G = \mathbb{Z}_5^2$ of $\mathbb{C}(S)$ over $\mathbb{C}(\mathbb{P}^1 \times \mathbb{P}^1)$ acts trivially on $\mathbb{C}(\Sigma)$. It follows that $\mathbb{C}(\Sigma) = \mathbb{C}(S)$, namely φ is birational.

Acknowledgements We are grateful to the referee for his help in improving the presentation and for suggesting the calculation in Example 4.4.

This work began during the final meeting in Torino of the EU HCM network AGE contract ERBCHRXT 940557. The research has been partly supported by the Italian P.I.N. 1997 Geometria algebrica, algebra commutativa e aspetti computazionali and by Project PRAXIS XXI 2/2.1/MAT/73/94. The first author is a member of CMAF and of the Departamento de Matemática da Faculdade de Ciências da Universidade de Lisboa and the second author is a member of GNSAGA of CNR.

References

- [1] A. BEAUVILLE, ‘Complex Algebraic Surfaces’, Cambridge University Press (1983).
- [2] M. BELTRAMETTI, A. SOMMESE, ‘Zero-cycles and k-th order embeddings of smooth projective surfaces’, in *Problems in the theory of surfaces and their classification* (F. Catanese, C. Ciliberto, M. Cornalba eds.), Symposia Math. 32 (1991), 33–48.
- [3] P. BURNIAT, ‘Sur les surfaces de genre $P_{12} > 0$ ’, *Ann. Mat. Pura Appl.*, (4) 71 (1966).
- [4] F. CATANESE, ‘Singular bidouble covers and the construction of interesting algebraic surfaces’ in *Algebraic Geometry: Hirzebruch 70*, A.M.S. Contemporary Mathematics, vol. 241 (1999).
- [5] F. CATANESE, C. CILIBERTO, ‘On the irregularity of cyclic coverings of algebraic varieties’, in *Geometry of Complex Projective Varieties - Cetraro*, (Italy), June 1990, (A. Lanteri, M. Palleschi, D. C. Struppa eds.), Mediterranean Press (1993), 89–116.
- [6] M. DE FRANCHIS, ‘Sugli integrali di Picard relativi a una superficie doppia’, *Rend. Circ. Mat. Palermo*, XX (1905), 331–334.
- [7] I. DOLGACHEV, ‘Algebraic surfaces with $p = p_g = 0$ ’, in *Algebraic surfaces* (C.I.M.E. 1977 Varenna), Liguori Napoli (1981), 97–215.
- [8] M. INOUE, ‘Some new surfaces of general type’, *Tokyo J. of Math.*, Vol. 17 No. 2 (1994), 295–319.

- [9] M. MENDES LOPES, ‘The degree of the generators of the canonical ring of surfaces of general type with $p_g = 0$ ’, *Arch. Math.*, 69 (1997), 435–440.
- [10] M. MENDES LOPES, R. PARDINI, ‘A connected component of the moduli space of surfaces with $p_g = 0$ ’, *Topology* (to appear); Eprint: math.AG/9910012.
- [11] M. NAGATA, ‘On rational surfaces I.’, *Mem. Coll. Sci., U. of Kyoto*, ser. A, vol. XXXII, Mathematics No. 3 (1960), 351–370.
- [12] R. PARDINI, ‘Abelian covers of algebraic varieties’, *J. reine angew. Math.* 417 (1991), 191–213.
- [13] C. PETERS, ‘On certain examples of surfaces with $p_g = 0$ due to Bur-
natiat’, *Nagoya Math. J.*, Vol. 166 (1977), 109–119.
- [14] I. REIDER, ‘Vector bundles of rank 2 and linear systems on algebraic
surfaces’, *Ann. of Math.*, 127 (1988), 309–316.
- [15] G. XIAO, ‘Finitude de l’application bicanonique des surfaces de type
général’, *Bull. Soc. Math. France*, 113 (1985), 23–51.
- [16] G. XIAO, ‘Degree of the bicanonical map of a surface of general type’,
Amer. J. of Math., 112 (5) (1990), 713–737.

Margarida Mendes Lopes
CMAF
Universidade de Lisboa
Av. Prof. Gama Pinto, 2
1649-003 Lisboa, PORTUGAL
mmlopes@lmc.fc.ul.pt

Rita Pardini
Dipartimento di Matematica
Università di Pisa
Via Buonarroti 2
56127 Pisa, ITALY
pardini@dm.unipi.it